

Dechant, Pierre-Philippe ORCID:

<https://orcid.org/0000-0002-4694-4010> (2016) A systematic construction of representations of quaternionic type. In: Alterman Conference on Geometric Algebra, 1st - 9th August 2016, Brasov, Romania. (Unpublished)

Downloaded from: <http://ray.yorks.ac.uk/id/eprint/4006/>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repository Policy Statement](#)

# RaY

Research at the University of York St John

For more information please contact RaY at [ray@yorks.ac.uk](mailto:ray@yorks.ac.uk)



# A systematic construction of representations of quaternionic type

Pierre-Philippe Dechant

Mathematics Department, University of York

Alterman Conference Brasov – August 4th, 2016

- 1 Polyhedral groups, Platonic solids and root systems
- 2 A Clifford way of doing orthogonal transformations
- 3 Clifford algebra and quaternions
- 4 Representations from multivector groups: representations of quaternionic type
- 5 Conclusions

# Platonic Solids



| Platonic Solid              | Group            | root system                   |
|-----------------------------|------------------|-------------------------------|
| Tetrahedron                 | $A_3$<br>$A_1^3$ | Cuboctahedron<br>Octahedron   |
| Octahedron<br>Cube          | $B_3$            | Cuboctahedron<br>+ Octahedron |
| Icosahedron<br>Dodecahedron | $H_3$            | Icosidodecahedron             |

- Platonic Solids have been known for millennia


# Platonic Solids

 $A_1^3$  $A_3$  $B_3$  $H_3$ 

| Platonic Solid                     | Group            | root system                   |
|------------------------------------|------------------|-------------------------------|
| <b>Tetrahedron</b>                 | $A_3$<br>$A_1^3$ | Cuboctahedron<br>Octahedron   |
| <b>Octahedron</b><br>Cube          | $B_3$            | Cuboctahedron<br>+ Octahedron |
| <b>Icosahedron</b><br>Dodecahedron | $H_3$            | Icosidodecahedron             |

- **Platonic Solids** have been known for millennia
- Described by **Coxeter** groups

# Root systems



reflection/Coxeter groups

**Root system**  $\Phi$ : set of vectors  $\alpha$  in a **vector space** with an **inner product** such that




$$1. \Phi \cap \mathbb{R}\alpha = \{-\alpha, \alpha\} \quad \forall \alpha \in \Phi$$

$$2. s_\alpha \Phi = \Phi \quad \forall \alpha \in \Phi$$

**Simple roots**: express every element of  $\Phi$  via a  **$\mathbb{Z}$ -linear combination**.



$$s_\alpha : v \rightarrow s_\alpha(v) = v - 2 \frac{(v|\alpha)}{(\alpha|\alpha)} \alpha$$

# The Icosahedron




- **Rotational** icosahedral group is  $I = A_5$  of order **60**
- **Full** icosahedral group is  $H_3$  of order **120** (including reflections/inversion); generated by the root system icosidodecahedron

# Non-crystallographic Coxeter groups $H_2 \subset H_3 \subset H_4$



$$A = \begin{pmatrix} 2 & -\tau \\ -\tau & 2 \end{pmatrix}$$



$$A = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -\tau \\ 0 & -\tau & 2 \end{pmatrix}$$



$$A = \begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -\tau \\ 0 & 0 & -\tau & 2 \end{pmatrix}$$

$H_2 \subset H_3 \subset H_4$ : 10, 120, 14,400 elements, the only Coxeter groups that generate **rotational symmetries of order 5** linear combinations now in the **extended integer ring**

$$\mathbb{Z}[\tau] = \{a + \tau b \mid a, b \in \mathbb{Z}\}$$

**golden ratio**

$$\tau = \frac{1}{2}(1 + \sqrt{5}) = 2 \cos \frac{\pi}{5}$$

$$x^2 = x + 1$$


$$\tau' = \sigma = \frac{1}{2}(1 - \sqrt{5}) = 2 \cos \frac{2\pi}{5}$$

$$\tau + \sigma = 1, \tau\sigma = -1$$



# Cartan-Dynkin diagrams

Coxeter-Dynkin diagrams: node = simple root, no link = roots orthogonal i.e. angle  $\frac{\pi}{2}$ , simple link = roots at angle  $\frac{\pi}{3}$ , link with label  $m = \text{angle } \frac{\pi}{m}$ .




# $H_3$ – the icosahedral group



$$\alpha_1 = (0, 1, 0), \quad \alpha_2 = -\frac{1}{2}(-\sigma, 1, \tau), \quad \alpha_3 = (0, 0, 1)$$

$$T_5 = (\tau, -1, 0), \quad T_3 = (\tau, 0, \sigma), \quad T_2 = (1, 0, 0)$$



Icosahedron, Dodecahedron, Icosidodecahedron ( $H_3$  root system)

- 1 Polyhedral groups, Platonic solids and root systems
- 2 A Clifford way of doing orthogonal transformations
- 3 Clifford algebra and quaternions
- 4 Representations from multivector groups: representations of quaternionic type
- 5 Conclusions

# Platonic Solids

 $A_1^3$  $A_1^4$  $A_3$  $D_4$  $B_3$  $F_4$  $H_3$  $H_4$ 


- Concatenating reflections gives **Clifford** spinors (**binary polyhedral groups**)

- These **induce 4D root systems**

$$R = a_0 + a_1 e_2 e_3 + a_2 e_3 e_1 + a_3 e_1 e_2 \Rightarrow$$

$$R\tilde{R} = a_0^2 + a_1^2 + a_2^2 + a_3^2$$

- 4D analogues of the Platonic Solids and give rise to 4D **Coxeter** groups



# Clifford Algebra and orthogonal transformations

- **Geometric Product** for two vectors  $ab \equiv a \cdot b + a \wedge b$
- **Inner product** is symmetric part  $a \cdot b = \frac{1}{2}(ab + ba)$
- Reflecting  $a$  in  $n$  is given by  $a' = a - 2(a \cdot n)n = -nan$  ( $n$  and  $-n$  **doubly cover** the same reflection)
- Via **Cartan-Dieudonné** theorem any orthogonal transformation can be written as **successive reflections**, which are **doubly covered** by Clifford versors/pinors  $A$

$$x' = \pm n_1 n_2 \dots n_k x n_k \dots n_2 n_1 =: \pm A x \tilde{A}$$

# Clifford Algebra of 3D

- E.g. **Pauli algebra** in 3D (likewise for **Dirac algebra** in 4D) is

$$\begin{array}{cccc}
 \underbrace{\{1\}} & \underbrace{\{e_1, e_2, e_3\}} & \underbrace{\{e_1 e_2, e_2 e_3, e_3 e_1\}} & \underbrace{\{I \equiv e_1 e_2 e_3\}} \\
 1 \text{ scalar} & 3 \text{ vectors} & 3 \text{ bivectors} & 1 \text{ trivector}
 \end{array}$$


- We can **multiply together root vectors** in this algebra  $\alpha_i \alpha_j \dots$
- A general element has **8** components, **even** products (rotations/spinors) have **four** components:

$$R = a_0 + a_1 e_2 e_3 + a_2 e_3 e_1 + a_3 e_1 e_2 \Rightarrow R \tilde{R} = a_0^2 + a_1^2 + a_2^2 + a_3^2$$

- So behaves as a **4D Euclidean** object – inner product

$$(R_1, R_2) = \frac{1}{2}(R_2 \tilde{R}_1 + R_1 \tilde{R}_2)$$

# Spinors from reflections



- The 6 **roots**  $(\pm 1, 0, 0)$  and permutations in  $A_1 \times A_1 \times A_1$  generate 8 **spinors**:
- $\boxed{\pm e_1, \pm e_2, \pm e_3}$  give the 8 spinors  $\boxed{\pm 1, \pm e_1 e_2, \pm e_2 e_3, \pm e_3 e_1}$
- This is a **discrete spinor group** isomorphic to the **quaternion** group  $Q$ .
- As 4D vectors these are  $(\pm 1, 0, 0, 0)$  and permutations, the 8 **roots** of  $A_1 \times A_1 \times A_1 \times A_1$  (the 16-cell).

# Induction Theorem – root systems

- Induction Theorem: Every rank-3 root system induces a rank-4 root system (and thereby Coxeter groups) via these 3D spinor groups.



# Induction Theorem – root systems

- Induction Theorem: Every rank-3 root system induces a rank-4 root system (and thereby Coxeter groups) via these 3D spinor groups.
- Check axioms:
  1.  $\Phi \cap \mathbb{R}\alpha = \{-\alpha, \alpha\} \quad \forall \alpha \in \Phi$
  2.  $s_\alpha \Phi = \Phi \quad \forall \alpha \in \Phi$

# Induction Theorem – root systems





- Induction Theorem: Every rank-3 root system induces a rank-4 root system (and thereby Coxeter groups) via these 3D spinor groups.
- Check axioms:
  1.  $\Phi \cap \mathbb{R}\alpha = \{-\alpha, \alpha\} \quad \forall \alpha \in \Phi$
  2.  $s_\alpha \Phi = \Phi \quad \forall \alpha \in \Phi$
- Proof: 1.  $R$  and  $-R$  are in a spinor group by construction (double cover of orthogonal transformations), 2. closure under reflections is guaranteed by the closure property of the spinor group (with a twist:  $-R_1 \tilde{R}_2 R_1$ )

# Spinors from reflections

- Symmetry groups of the **Platonic Solids**:
- The 6/12/18/30 **reflections** in  $A_1 \times A_1 \times A_1 / A_3 / B_3 / H_3$  generate 8/24/48/120 **spinors**.
- The **discrete spinor group** is isomorphic to the **quaternion group**  $Q$  / **binary tetrahedral group**  $2T$  / **binary octahedral group**  $2O$  / **binary icosahedral group**  $2I$ ).

# Spinors and Polytopes

- Can reinterpret **spinors in  $\mathbb{R}^3$**  as **vectors in  $\mathbb{R}^4$**
- Give (exceptional) root systems ( $D_4, F_4, H_4$ )
- They constitute the **vertices** of the **16-cell**, **24-cell**, **24-cell** and **dual 24-cell** and the **600-cell**
- These are 4D analogues of the **Platonic Solids**. **Strange symmetries** better understood in terms of **3D spinors**



# Root systems in three and four dimensions

The **spinors** from the reflections in the **rank-3 Coxeter group** via the geometric product are the **binary polyhedral groups**  $Q$ ,  $2T$ ,  $2O$  and  $2I$ , which generate (mostly exceptional) **rank-4 groups**, but **not known why**, and why the 'mysterious symmetries'.

| rank-3 group                | diagram | binary | rank-4 group                           | diagram |
|-----------------------------|---------|--------|----------------------------------------|---------|
| $A_1 \times A_1 \times A_1$ |         | $Q$    | $A_1 \times A_1 \times A_1 \times A_1$ |         |
| $A_3$                       |         | $2T$   | $D_4$                                  |         |
| $B_3$                       |         | $2O$   | $F_4$                                  |         |
| $H_3$                       |         | $2I$   | $H_4$                                  |         |

- 1 Polyhedral groups, Platonic solids and root systems
- 2 A Clifford way of doing orthogonal transformations
- 3 Clifford algebra and quaternions**
- 4 Representations from multivector groups: representations of quaternionic type
- 5 Conclusions

# Quaternion groups via the geometric product

- The 8 quaternions of the form  $(\pm 1, 0, 0, 0)$  and permutations are the **Lipschitz units**, the **quaternion group** in 8 elements.
- The 8 Lipschitz units together with  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  are the **Hurwitz units**, the **binary tetrahedral group** of order 24.  
Together with the 24 'dual' quaternions of the form  $\frac{1}{\sqrt{2}}(\pm 1, \pm 1, 0, 0)$ , they form the **binary octahedral group** of order 48.
- The 24 Hurwitz units together with the 96 unit quaternions of the form  $(0, \pm \tau, \pm 1, \pm \sigma)$  and even permutations, are called the **Icosians**. The icosian group is isomorphic to the **binary icosahedral group** with 120 elements.
- The unit spinors  $\{1; e_2 e_3; e_3 e_1; e_1 e_2\}$  of  $\text{Cl}(3)$  are isomorphic to the **quaternion algebra**  $\mathbb{H}$ .

## $H_4$ from icosahedral spinors

- The  $H_3$  root system has 30 **roots** e.g. simple roots  $\alpha_1 = e_2$ ,  $\alpha_2 = -\frac{1}{2}((\tau-1)e_1 + e_2 + \tau e_3)$  and  $\alpha_3 = e_3$ .
- The subgroup of **rotations** is  $A_5$  of order **60**
- These are doubly covered by **120** spinors of the form  
 $\alpha_1 \alpha_2 = -\frac{1}{2}(1 - (\tau-1)e_1 e_2 + \tau e_2 e_3)$ ,  $\alpha_1 \alpha_3 = e_2 e_3$  and  
 $\alpha_2 \alpha_3 = -\frac{1}{2}(\tau - (\tau-1)e_3 e_1 + e_2 e_3)$ .
- As a set of **vectors** in 4D, they are

$(\pm 1, 0, 0, 0)$  (8 permutations) ,  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  (16 permutations)

$\frac{1}{2}(0, \pm 1, \pm \sigma, \pm \tau)$  (96 even permutations) ,

which are precisely the 120 roots of the  **$H_4$  root system**.



# Systematic construction of the polyhedral groups


- Multiplying together root vectors in the Clifford algebra gave a **systematic** way of constructing the **binary polyhedral** groups as 3D spinors = **quaternions**.
- The 6/12/18/30 **roots** in  $A_1 \times A_1 \times A_1 / A_3 / B_3 / H_3$  generate 8/24/48/120 **spinors**.
- The **discrete spinor group** is isomorphic to the **quaternion** group  $Q$  / **binary tetrahedral** group  $2T$  / **binary octahedral** group  $2O$  / **binary icosahedral** group  $2I$ ).

|         |       |       |       |
|---------|-------|-------|-------|
| $A_1^3$ | $A_3$ | $B_3$ | $H_3$ |
| $A_1^4$ | $D_4$ | $F_4$ | $H_4$ |

# Quaternionic representations of 3D and 4D Coxeter groups

- Groups  $E_8$ ,  $D_4$ ,  $F_4$  and  $H_4$  have representations in terms of **quaternions**
- **Extensively used** in the high energy physics/quasicrystal/Coxeter/polytope literature and thought of as deeply significant, though not really clear why
- e.g.  $H_4$  consists of 120 elements of the form  $(\pm 1, 0, 0, 0)$ ,  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  and  $(0, \pm \tau, \pm 1, \pm \sigma)$
- Seen as remarkable that the **subset of the 30 pure quaternions** is a realisation of  $H_3$  (**a sub-root system**)
- Similarly,  $B_3$  and  $A_1 \times A_1 \times A_1$  have representations in terms of **pure quaternions**
- Clifford provides a **much simpler geometric explanation**

# Quaternionic representations in the literature



Pure quaternions = Hodge dualised **root vectors**

Quaternions = **spinors**

# Demystifying Quaternionic Representations

- **Pure quaternion subset** of 4D groups only gives 3D group if the 3D group **contains the inversion/pseudoscalar  $I$**
- e.g. **does not work** for the tetrahedral group  $A_3$ , but  $A_3 \rightarrow D_4$  **induction still works**, with the central node essentially 'spinorial'
- In fact, it goes the other way around: the **3D groups induce the 4D groups** via spinors
- The rank-4 groups are also generated (under quaternion multiplication) by two quaternions we can identify as  **$R_1 = \alpha_1 \alpha_2$  and  $R_2 = \alpha_2 \alpha_3$**
- Can see these are '**spinor generators**' and how they don't really contain any more information/roots than the rank-3 groups alone

# Quaternions vs Clifford versions

- **Sandwiching** is often seen as particularly nice feature of the quaternions giving rotations
- This is actually a **general feature** of Clifford algebras/versions in any dimension; the isomorphism to the quaternions is **accidental** to 3D
- However, the **root system** construction does not necessarily generalise
- 2D generalisation merely gives that  $I_2(n)$  is **self-dual**
- **Octonionic** generalisation just induces two copies of the above 4D root systems, e.g.  $A_3 \rightarrow D_4 \oplus D_4$
- Recently constructed  $E_8$  from the 240 pinors doubly covering 120 elements of  $H_3$  in  $2^3 = 8$ -dimensional 3D Clifford algebra

- 1 Polyhedral groups, Platonic solids and root systems
- 2 A Clifford way of doing orthogonal transformations
- 3 Clifford algebra and quaternions
- 4 Representations from multivector groups: representations of quaternionic type**
- 5 Conclusions

# Polyhedral groups as multivector groups

| Group     | Discrete subgroup         | Order  | Action Mechanism                 |
|-----------|---------------------------|--------|----------------------------------|
| $SO(3)$   | rotational (chiral)       | $ G $  | $x \rightarrow \tilde{R}xR$      |
| $O(3)$    | reflection (full/Coxeter) | $2 G $ | $x \rightarrow \pm \tilde{A}xA$  |
| $Spin(3)$ | binary                    | $2 G $ | $(R_1, R_2) \rightarrow R_1 R_2$ |
| $Pin(3)$  | pinory (?)                | $4 G $ | $(A_1, A_2) \rightarrow A_1 A_2$ |

- e.g. the **chiral icosahedral** group has 60 elements, encoded in GA by 120 rotors, which form the **binary icosahedral** group
- together with the **inversion/pseudoscalar**  $I$  this gives 60 rotations and 60 rotoinversions, i.e. the **full icosahedral** group  $H_3$  in 120 elements doubly covered by 240 pinors

# Some Group Theory: chiral, full, binary, pin

- Easy to calculate **conjugacy classes** etc of versors in GA
- Chiral (**binary**) polyhedral groups have irreps
- tetrahedral (12/**24**): 1, 1', 1'',  $2_s$ ,  $2'_s$ ,  $2''_s$ , 3
- octahedral (24/**48**): 1, 1', 2,  $2_s$ ,  $2'_s$ , 3, 3',  $4_s$
- icosahedral (60/**120**): 1,  $2_s$ ,  $2'_s$ , 3,  $\bar{3}$ , 4,  $4_s$ , 5,  $6_s$
- All binary are **discrete subgroups of  $SU(2)$**  and all thus have a  $2_s$  spinor irrep
- Connection with **Trinities and the McKay correspondence**



# Representations from Clifford multivector groups

- The usual picture of **orthogonal transformations** on an  $n$ -dimensional vector space is via  $n \times n$  **matrices** acting on vectors, immediately making connections with **representations = matrices satisfying the group multiplication laws**.
- **Easy to construct representations** with (s)pinors in the  $2^n$ -dimensional Clifford algebra as **reshuffling components**.
- Spinors leave the **original**  $n$ -dimensional **vector** space invariant, **reshuffle** the components of the **vector**.
- But can also consider various representation matrices acting on **different subspaces** of the Clifford algebra.

# Representations from Clifford multivector groups – trivial, parity, rotation representations

- The **scalar** subspace is **one-dimensional**.  $\tilde{R}1R = \tilde{R}R = 1$  gives the **trivial representation**, and likewise pinors  $A$  give the **parity**.
- The double-sided action  $\tilde{R}xR$  of spinors  $R$  on a **vector**  $x$  in the  $n$ -dimensional vector space gives an  $n \times n$ -dimensional representation, which is just the usual **rotation matrices**.
- E.g.  $e_1e_2$  acting on  $x = x_1e_1 + x_2e_2 + x_3e_3$  gives  $e_2e_1xe_1e_2 = -x_1e_1 - x_2e_2 + x_3e_3$  which could also be expressed as 
$$\begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -x_1 \\ -x_2 \\ x_3 \end{pmatrix}$$
- If the spinors were acting as  $Rx\tilde{R}$  would give a **potentially different representation**.

# Characters, their norm, and the Frobenius-Schur indicator

- **Similarity** transformed representations are also good representations, but are not fundamentally different: they are **equivalent**.
- So want a measure for a representation that is **invariant** under similarity transformations, e.g. the **trace** aka the **character**  $\chi$  of a matrix
- A **class function** i.e. the same within a conjugacy class because of the cyclicity of the trace
- The **character norm**  $||\chi||^2 := \frac{1}{|G|} \sum_{g \in G} |\chi(g)|^2$
- The **Frobenius-Schur indicator**  $\nu := \frac{1}{|G|} \sum_{g \in G} \chi(g^2)$

# Real representations of real, complex, and quaternionic type

- $||\chi||^2 = \frac{1}{|G|} \sum_{g \in G} |\chi(g)|^2 = 1$ : representation of **real** type
- $||\chi||^2 = \frac{1}{|G|} \sum_{g \in G} |\chi(g)|^2 = 2$ : representation of **complex** type
- $||\chi||^2 = \frac{1}{|G|} \sum_{g \in G} |\chi(g)|^2 = 4$ : representation of **quaternionic** type
- Theorem: A complex representation is irreducible if and only if  $||\chi||^2 = 1$ .
- Theorem: A **real** representation is **irreducible** if and only if  $||\chi||^2 + \nu(\chi) = 2$ , e.g.  $4 - 2 = 2$  or  $1 + 1 = 2$ .

# Representations from Clifford multivector groups – $8 \times 8$ and $4 \times 4$ (whole algebra / even subalgebra)

- Rather than restricting oneself to the  $n$ -dimensional vector space, one can also define representations by  $2^n \times 2^n$ -matrices acting on the **whole** Clifford algebra, i.e. any element acting on an arbitrary element, e.g. here  $8 \times 8$ .
- Likewise, one can define  $2^{(n-1)} \times 2^{(n-1)}$ -dimensional spinor representations as acting on the **even subalgebra**.
- 3D spinors have **components** in  $(1, e_1 e_2, e_2 e_3, e_3 e_1)$ , **multiplication** with another spinor e.g.  $e_1 e_2$  will **reshuffle** these components  $(e_1 e_2, -1, -e_3 e_1, e_2 e_3)$
- This **reshuffling** can therefore be described by a  $4 \times 4$ -matrix.

# 4 × 4 – explicit example: $A_1^3$

- E.g.  $\boxed{\pm e_1, \pm e_2, \pm e_3}$  give the 8 spinors  
 $\boxed{\pm 1, \pm e_1 e_2, \pm e_2 e_3, \pm e_3 e_1}$ , or  $(\pm 1, 0, 0, 0)$  (8 permutations)
- $\|\chi\|^2 = 32/8 = 4$ ,  $v = -2$  and  $\|\chi\|^2 + v = 2$  i.e. **real**  
**irreducible of quaternionic type**

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, \\
 \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}$$

# Character table of $Q$

| $Q$   | 1 | -1 | $\pm e_1 e_2$ | $\pm e_2 e_3$ | $\pm e_3 e_1$ |
|-------|---|----|---------------|---------------|---------------|
| 1     | 1 | 1  | 1             | 1             | 1             |
| 1'    | 1 | 1  | -1            | -1            | 1             |
| 1''   | 1 | 1  | -1            | 1             | -1            |
| 1'''  | 1 | 1  | 1             | -1            | -1            |
| $4_H$ | 4 | -4 | 0             | 0             | 0             |

## $4 \times 4$ – explicit example: $A_3$

- As a set of **vectors** in 4D, they are  $(\pm 1, 0, 0, 0)$  (8 permutations),  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  (16 permutations)
- Conjugacy classes:  
 $1 \cdot 4^2 + 1 \cdot (-4)^2 + 6 \cdot 0^2 + 8 \cdot 2^2 + 8 \cdot (-2)^2 = 32 + 32 + 32 = 96$
- $\|\chi\|^2 = 96/24 = 4$ ,  $\nu = -2$  and  $\|\chi\|^2 + \nu = 2$  i.e. **real irreducible of quaternionic type**.



## $3 \times 3$ – explicit example: $H_3$

- Icosahedral spinors are

$(\pm 1, 0, 0, 0)$  (8 permutations),  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  (16 permutations)

$\frac{1}{2}(0, \pm 1, \pm \sigma, \pm \tau)$  (96 even permutations),

- E.g. the rotation matrices corresponding to  $\alpha_1 \alpha_2$  and  $\alpha_2 \alpha_3$  via  $\tilde{R}XR$  are

$$\frac{1}{2} \begin{pmatrix} \tau & \tau - 1 & -1 \\ 1 - \tau & -1 & -\tau \\ -1 & \tau & 1 - \tau \end{pmatrix} \text{ and } \frac{1}{2} \begin{pmatrix} \tau & 1 - \tau & -1 \\ 1 - \tau & 1 & -\tau \\ 1 & \tau & \tau - 1 \end{pmatrix}.$$

The characters  $\chi(g)$  are obviously 0 and  $\tau$

- $||\chi||^2 = 120/120 = 1$ ,  $\nu = 1$  and  $||\chi||^2 + \nu = 2$  i.e. **real irreducible of real type**

## $3 \times 3$ – explicit example: $H_3$ other way

- If the spinors were acting as  $R \times \tilde{R}$ , then

$$\frac{1}{2} \begin{pmatrix} \tau & 1-\tau & -1 \\ \tau-1 & -1 & \tau \\ -1 & -\tau & 1-\tau \end{pmatrix} \text{ and } \frac{1}{2} \begin{pmatrix} \tau & 1-\tau & 1 \\ 1-\tau & 1 & \tau \\ -1 & -\tau & \tau-1 \end{pmatrix},$$

with the same characters as before. Swapping the action of the spinor can change the representation.

## $4 \times 4$ – explicit example: $H_3$

- Spinors  $\alpha_1 \alpha_2$  and  $\alpha_2 \alpha_3$  multiplying a generic spinor  $R = a_0 + a_1 e_2 e_3 + a_2 e_3 e_1 + a_3 e_1 e_2$  from the left reshuffles the components  $(a_1, a_2, a_3, a_0)$  with the matrices given as

$$\frac{1}{2} \begin{pmatrix} -1 & \tau - 1 & 0 & -\tau \\ 1 - \tau & -1 & -\tau & 0 \\ 0 & \tau & -1 & \tau - 1 \\ \tau & 0 & 1 - \tau & -1 \end{pmatrix}, \frac{1}{2} \begin{pmatrix} -\tau & 0 & 1 - \tau & -1 \\ 0 & -\tau & -1 & \tau - 1 \\ \tau - 1 & 1 & -\tau & 0 \\ 1 & 1 - \tau & 0 & -\tau \end{pmatrix},$$

with characters  $-2$  and  $-2\tau$ .

## $4 \times 4$ – explicit example $H_3$ : quaternionic type

- 120  $4 \times 4$  matrices – 9 conjugacy classes, with pairs that have  $\pm 2\chi_3$  so gives **4 times** that of the  $3 \times 3$  case
- $|G| \cdot \|\chi\|^2 = 1 \cdot 4^2 + 1 \cdot (-4)^2 + 12 \cdot (-2\tau)^2 + 12 \cdot (2\tau)^2 + 12 \cdot (-2\sigma)^2 + 12 \cdot (2\sigma)^2 + 20 \cdot (-2)^2 + 20 \cdot (2)^2 + 30 \cdot 0^2 = \mathbf{480}$
- $\|\chi\|^2 = 480/120 = \mathbf{4}$ ,  $\nu = \mathbf{-2}$  and  $\|\chi\|^2 + \nu = \mathbf{2}$  i.e. **real irreducible of quaternionic type**

# Character table of $I = A_5$

| $I$       | 1 | $20C_3$ | $15C_2$ | $12C_5$  | $12C_5^2$ |
|-----------|---|---------|---------|----------|-----------|
| 1         | 1 | 1       | 1       | 1        | 1         |
| 3         | 3 | 0       | -1      | $\tau$   | $\sigma$  |
| $\bar{3}$ | 3 | 0       | -1      | $\sigma$ | $\tau$    |
| 4         | 4 | 1       | 0       | -1       | -1        |
| 5         | 5 | -1      | 1       | 0        | 0         |

# Character table of $2I$

| $I$             | 1 | $20C_3$ | $30C_2$ | $12C_5$    | $12C_5^2$  | $-1$ | $-20C_3$ | $-12C_5$  | $-12C_5^2$ |
|-----------------|---|---------|---------|------------|------------|------|----------|-----------|------------|
| 1               | 1 | 1       | 1       | 1          | 1          | 1    | 1        | 1         | 1          |
| 3               | 3 | 0       | -1      | $\tau$     | $\sigma$   | 3    | 0        | $\tau$    | $\sigma$   |
| $\bar{3}$       | 3 | 0       | -1      | $\sigma$   | $\tau$     | 3    | 0        | $\sigma$  | $\tau$     |
| 4               | 4 | 1       | 0       | -1         | -1         | 4    | 1        | -1        | -1         |
| 5               | 5 | -1      | 1       | 0          | 0          | 5    | -1       | 0         | 0          |
| 2               | 2 | -1      | 0       | $-\sigma$  | $-\tau$    | -2   | 1        | $\sigma$  | $\tau$     |
| 2               | 2 | -1      | 0       | $-\tau$    | $-\sigma$  | -2   | 1        | $\tau$    | $\sigma$   |
| 4               | 4 | 1       | 0       | -1         | -1         | -4   | -1       | 1         | 1          |
| 6               | 6 | 0       | 0       | 1          | 1          | -6   | 0        | -1        | -1         |
| $4_H$           | 4 | -2      | 0       | $-2\tau$   | $-2\sigma$ | -4   | 2        | $2\tau$   | $2\sigma$  |
| $4_{\tilde{H}}$ | 4 | -2      | 0       | $-2\sigma$ | $-2\tau$   | -4   | 2        | $2\sigma$ | $2\tau$    |

# A general construction of representations of quaternionic type – canonical representations

- It had so far been **overlooked** that there is a **systematic construction** of representations of **quaternionic type** for 3D polyhedral groups
- This is simply due to the fact that the **spinors** in 3D provide a realisation of the **quaternions**
- Therefore spinors provide 4x4 representations of quaternionic type for **all** (though limited number of) possible groups
- However, they are **canonical** for a choice of 3D **simple roots**, i.e. there is a preferred amongst all similarity transformed versions
- These **simple roots** also determine the 3x3 **rotation** matrices and their **reversed** representations in a similar **canonical** way

# Characters in general

- For a **general spinor**  $R = a_0 + a_1 e_2 e_3 + a_2 e_3 e_1 + a_3 e_1 e_2$  one has **3D character**  $\chi = 3a_0^2 - a_1^2 - a_2^2 - a_3^2$  and **representation**

$$\frac{1}{2} \begin{pmatrix} a_0^2 + a_1^2 - a_2^2 - a_3^2 & -2a_0 a_3 + 2a_1 a_2 & 2a_0 a_2 + 2a_1 a_3 \\ 2a_0 a_3 + 2a_1 a_2 & a_0^2 - a_1^2 + a_2^2 - a_3^2 & -2a_0 a_1 + 2a_2 a_3 \\ -2a_0 a_2 + 2a_1 a_3 & 2a_0 a_1 + 2a_2 a_3 & a_0^2 - a_1^2 - a_2^2 + a_3^2 \end{pmatrix}$$

- and the **4D rep and character** are

$$\begin{pmatrix} a_0 & a_3 & -a_2 & a_1 \\ -a_3 & a_0 & a_1 & a_2 \\ a_2 & -a_1 & a_0 & a_3 \\ -a_1 & -a_2 & -a_3 & a_0 \end{pmatrix}, \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_0 \end{pmatrix} \text{ and } \chi = 4a_0.$$

- Characters** of the representations are **all** determined by the **spinor**!



- 1 Polyhedral groups, Platonic solids and root systems
- 2 A Clifford way of doing orthogonal transformations
- 3 Clifford algebra and quaternions
- 4 Representations from multivector groups: representations of quaternionic type
- 5 Conclusions**

# Conclusions

- **General construction** of 4D root systems from 3D root systems – connections with **McKay correspondence**, **Trinities**, **Moonshine** etc
- Construction **systematically** and **canonically** gives representations of 4D root systems and 3D root systems in terms of **(pure) quaternions**
- Construction **systematically** and **canonically** gives construction of the polyhedral groups and their representations, in particular trivial, rotation and spinor representations of **quaternionic type** with relations among them and their characters

# Arnold's Trinities

- **Arnold** noticed that often **real**, **complex** and **quaternionic** versions of a theory are remarkably similar
- **Trinities**  $(\mathbb{R}, \mathbb{C}, \mathbb{H})$
- $(\mathbb{R}P^n, \mathbb{C}P^n, \mathbb{H}P^n)$ ,  $(\mathbb{R}P^1 = S^1, \mathbb{C}P^2 = S^2, \mathbb{H}P^1 = S^4)$ , the Möbius/Hopf bundles  $(S^1 \rightarrow S^1, S^4 \rightarrow S^2, S^7 \rightarrow S^4)$ ,  $(E_6, E_7, E_8)$
- **New connection** between  $(A_3, B_3, H_3)$  and  $(D_4, F_4, H_4)$  (and  $(E_6, E_7, E_8)$ !) via my **Clifford spinor construction**
- Also  $(24, 48, 120)$ , binary polyhedral groups  $(2T, 2O, 2I)$  and  $(12, 18, 30)$  (see McKay correspondence)

# The McKay Correspondence

binary polyhedral groups  
 $2T, 2O, 2I$   
 $\sum d_i$  12, 18, 30  
 $\sum d_i^2$  24, 48, 120

McKay correspondence


Exceptional  
Lie Groups

$E_6$ , 12


$E_7$ , 18

$E_8$ , 30

(Coxeter numbers)



# The McKay Correspondence



# Some Group Theory: chiral, full, binary, pin

- Easy to calculate **conjugacy classes** etc of versors in GA
- Chiral (**binary**) polyhedral groups have irreps
- tetrahedral (12/24): 1, 1', 1'',  $2_s$ ,  $2'_s$ ,  $2''_s$ , 3
- octahedral (24/48): 1, 1', 2,  $2_s$ ,  $2'_s$ , 3, 3',  $4_s$
- icosahedral (60/120): 1,  $2_s$ ,  $2'_s$ , 3,  $\bar{3}$ , 4,  $4_s$ , 5,  $6_s$
- All binary are **discrete subgroups of  $SU(2)$**  and all thus have a  $2_s$  spinor irrep
- Connection with **Trinities and the McKay correspondence**

# The McKay Correspondence

More than E-type groups: the infinite family of 2D groups, the **cyclic** and **dicyclic groups** are in correspondence with  $A_n$  and  $D_n$ , e.g. the quaternion group  $Q$  and  $D_4^+$ . So McKay correspondence not just a trinity but **ADE-classification**. We also have  $I_2(n)$  on top of the trinity ( $A_3, B_3, H_3$ )

| rank-3 group                | diagram | binary | rank-4 group                           | diagram | Lie algebra | diagram |
|-----------------------------|---------|--------|----------------------------------------|---------|-------------|---------|
| $A_1 \times A_1 \times A_1$ |         | $Q$    | $A_1 \times A_1 \times A_1 \times A_1$ |         | $D_4^+$     |         |
| $A_3$                       |         | $2T$   | $D_4$                                  |         | $E_6^+$     |         |
| $B_3$                       |         | $2O$   | $F_4$                                  |         | $E_7^+$     |         |
| $H_3$                       |         | $2I$   | $H_4$                                  |         | $E_8^+$     |         |